

Bioinformatics I: Protein Structure

Lecture 3 — Aims

- To introduce some of the current challenges in structural bioinformatics.

Lecture 1 — Objectives

After this lecture you will:

- understand the objectives of comparative modelling, fold recognition and secondary structure prediction.
- be aware of approaches to comparative modelling, fold recognition and secondary structure prediction.

Comparative modelling strategy

- identify a known structure that is predicted to be similar;
- align sequences;
- predict structurally conserved regions, and locations of insertions and deletions (sometimes called “indels”);
- build model backbone structure
 - copy predicted conserved main chain regions from template structure,
 - remodel loops with insertions or deletions;
- add side chains to the modelled main chain;
- evaluate and refine model.

Why build model structures?

Knowledge of a protein's three-dimensional structure is vital to a full understanding of the molecular basis for its biological function.

We want to understand the function of all proteins encoded by a genome, therefore we would like to know all of their 3-D structures.

Experimental techniques for determining protein structure are relatively slow and expensive, so we look to modelling as a way of extending the set of 3-D structures.

Modelling can also be used in protein engineering when designing proteins for therapeutic applications.

Energy calculations

Terms used in evaluating the energy of a conformation typically include:

- bond stretching
- bond angle bend
- terms penalising deviation from planarity, etc.
- torsion angles
- Van der Waals interactions
- hydrogen bonds
- electrostatics
- interactions with solvent, water and cosolutes

Fold recognition

The idea behind “threading”:

Imagine a wire wound into the shape of a known protein's main chain “fold”.

Imagine next that our new sequence is represented by beads that are “threaded”, in order, onto the wire, and are pushed along the wire.

At each step, a score is calculated based on which residues are adjacent in space, which residues are buried, etc.

Repeat this process for each different known fold.

A high score indicates that the sequence is compatible with that fold.

Heuristics for manual secondary structure prediction

- Many α -helices are amphipathic. Conserved hydrophobic residues at positions i , $i+3$, $i+4$, $i+7$, etc. are highly indicative of an α -helix.
- Half-buried strands will tend to have hydrophobic and hydrophilic residues at alternate positions.
- In proteins containing both α -helices and strands the strands are often completely buried and tend to contain only hydrophobic residues.

For more details and references, see:

<http://www.bmm.icnet.uk/people/rob/CCP11BBS/secstrucpred.html>

Secondary structure prediction

If neither sequence comparison nor fold recognition identifies a structure that can be used as a template for comparative modelling, then we can consider predicting secondary structure elements and how these might be assembled into a compact structure.

However, as noted by Ponder and Richards (1987):

“a major problem lies in the secondary structure prediction itself ... the problem appears to lie in the non-negligible effect of long-range tertiary structural features upon secondary structure”

and

“the problem of docking the preformed secondary units is formidable when considered in atomic detail.”

Alternative secondary structure assignment methods

Cuff J. A. and Barton G. J. Evaluation and improvement of multiple sequence methods for protein secondary structure prediction, *PROTEINS: Structure, Function and Genetics*. 34:508-519 (1999)

“Secondary structure definition methods DSSP[38], DEFINE[39] and STRIDE[40] were compared. All three agree at only 75% of positions. This is mainly due to differences between DEFINE and DSSP/STRIDE. DSSP and STRIDE agree at 95% of positions, though DSSP defines many more 4 residue helices than STRIDE.”

[38] W. Kabsch and C. Sander. A dictionary of protein secondary structure. *Biopolymers*, 22:2577-2637, 1983.

[39] F. M. Richards and C. E. Kundrot. Identification of structural motifs from protein coordinate data: secondary structure and first-level supersecondary structure. *Proteins*, 3:71-84, 1988.

[40] D. Frishman and P. Argos. Knowledge-based protein secondary structure assignment. *Proteins*, 23:566-579, 1995.